














Ek 3- Şekli Anahtar Örneği
















Major groups

ILLUSTRATED KEY

KEY TO MAJOR GROUPS

1	Plants reproducing by spores		Group A: Pteridophytes
	Plants reproducing by cones		<i>Ephedra foliata</i> (Gymnosperm)
	Plants reproducing by flowers		2
2	Submerged (marine or freshwater) plants		3
	Terrestrial or epiphytic plants (sometimes rooted in water but never submerged)		4
3	Plants submerged in fresh or brackish water		Group B: Fresh or brackish water monocots
	Plants submerged in the sea		Group C: Marine monocots
4	Leaves with parallel veins; flowers with parts in 3s		Group D: Terrestrial monocots
	Leaves with pinnate veins; flowers with parts in 2s, 4s or 5s, or parts many (dicots)		5
5	Annual or perennial herbs, sometimes woody at base		Group E: Herbaceous dicots
	Shrubs or trees		6
6	Leaves opposite		Group F: Woody dicots with opposite leaves
	Leaves alternate		Group G: Woody dicots with alternate leaves

GROUP A: PTERIDOPHYTES

1	Fronds simple or divided into lobes or dichotomously divided into linear segments		2		
	Fronds 1-pinnate		3		
	Fronds >1-pinnate		4		
2	Fronds ovate to elliptic with a cuneate base		Sporangia borne in two rows in a separate spike		<i>Diplazium polyphyllum</i> (Ophioglossaceae)
	Fronds simple, broadly ovate with a cordate or trincate base		Sporangia as above		<i>Diplazium reticulatum</i> (Ophioglossaceae)
	Fronds simple,		Sporangia borne (gathered) on narrow elliptic, elongate sori		<i>Loxogramme alpestrica</i> (Polypodiaceae)
	Fronds simple, narrowly elliptic		Sporangia borne in round sori on the underside of the frond		<i>Pteris macrocarpa</i> (Polypodiaceae)
	Fronds divided into 4 obovate leaflets		Sporangia borne in specialized pinnae (sporocarps)		<i>Marsilea coccinea</i> (Marsiliaceae)
	Fronds dichotomously divided into linear segments		Sporangia enclosed in reflexed frond margins		<i>Adiantum semibarbatum</i> (Adiantaceae)